

Gran-I-Grit Products

The Pioneer of Large Grit Feeding

GRAN-I-GRIT makes money for *you!*

When fed in the proper sizes, GRAN-I-GRIT usually pays for itself many times over by giving more broiler and turkey meat per pound of feed even on all mash rations. Layers fed GRAN-I-GRIT produce up to 20% more eggs with the same amount of feed.

How does GRAN-I-GRIT make this possible?

Birds rush feed through their digestive tracts, keeping it only from 1/2 to 12 minutes in the gizzard. The proper sizes of GRAN-I-GRIT in the gizzard grinds feed particles so small that the bird's digestive juices can quickly act on every bit of the valuable proteins, carbohydrates, fats, minerals and vitamins locked within the feed particles, converting them into a form for absorption into the blood stream where they aid growth and egg production. So you can see – without grit much of the valuable feed passes right through the bird *unused*.



151 Granite Quarry Trail
P. O. Box 151
Mount Airy, NC 27030
www.ncgranite.com

The North Carolina Granite Corporation



The North Carolina Granite Corporation

THE WORLD'S LARGEST OPEN FACE GRANITE QUARRY

GRAN-I-GRIT

Starter

Grower

Developer-Layer

Turkey

Turkey Finisher



GRAN-I-GRIT

Rough - Tough - Insoluble

The rough and tough surfaces of GRAN-I-GRIT enable the gizzard to cut up shavings, feathers and other fibrous materials that can clog up the digestive tract.

Attractive

The whiteness of GRAN-I-GRIT with contrasting black sparkling mica attracts birds to eat all they require.

Preventive

Continuous feeding of GRAN-I-GRIT also helps prevent digestive disturbances, diarrhea, cannibalism, cropbound conditions, and certain forms of paralysis. After passing through the digestive tract, GRAN-I-GRIT exercises and strengthens the organs, removing mucus and lessening the chance of disease germs being established.

Broilers: Feed Starter size for two weeks. Feed Grower size for next four weeks. Feed developer-Layer size after six weeks.

Layers: Feed Starter size for three weeks. Feed grower size for next four weeks. Feed Developer-Layer size after seven weeks.

Turkeys: Feed Grower for size for three weeks. Feed Developer-Layer size for next five weeks. Feed Turkey size for next six weeks. Feed Turkey finisher size after fourteenth week.



"STONEMO"
The Sturdy Bird

Amount to Feed

Usually a bird will eat no more grit than it needs, which for a layer is 1/6 to 1/3 lbs per month and for an adult turkey 1/3 to 2/3 lbs per month. If an occasional flock eats too much grit, change to the next larger size.

Birds pass out grit that is too small. The best rule is to feed the largest size the birds will eat in normal quantities.

Caution: Baby chicks and poults can mistake grit for feed with damaging results. Help them establish correct feeding habits by sprinkling grit over their mash or grain for the first two days. Then feed GRAN-I-GRIT in separate hoppers.

Gran-I-Grit Products

Starter
1/16" - 3/32"



Grower
3/32" - 3/16"



Developer - Layer
3/16" - 5/16"



Turkey
5/16" - 7/16"



Turkey Finisher
7/16" - 5/8"



**** All of our GRAN-I-GRIT products come ready for pick-up in 50 lb. poly bags with a 40 bag pallet minimum.**



Distributors:

Cooperative Feed Dealers Inc.

P.O. Box 670
Conklin, NY 13748-0670
1 800 333-0895

Mid-Central Products Inc.

P.O. Box 50068
Indianapolis, IN 46250-0068
317 570-7475

**** Or contact us directly.**

Direct 336 786-5141
Toll Free 1 800 227-6242
Email: grit@ncgranite.com

The North Carolina Granite Corporation

151 Granite Quarry Trail
P. O. Box 151
Mount Airy, NC 27030
www.ncgranite.com

Phone: 336-786-5141
Fax: 336-719-2623
grit@ncgranite.com